



1040-6182(95)00049-6

## GLOBAL YOUNGER DRYAS?

D. Peteet

NASA/Goddard Institute for Space Studies, 2880 Broadway, NY, NY 10025 and Lamont Doherty Earth Observatory, Palisades, NY 10964, U.S.A.

### INTRODUCTION

#### *Objectives of the Working Group*

The existence of the Younger Dryas cold event has been known for over a century. First named and described from buried leaves and fruits of *Dryas octopetala* (Rosaceae family) in Scandinavia, the re-appearance of these tundra indicators after shrubs and trees had begun to colonize a glaciated landscape (the Allerød warming), suggested a reversion to colder conditions (Jensen, 1938). Numerous stratigraphic studies subsequently proved the cooling as characteristic of late-glacial pollen diagrams throughout Europe 11–10,000  $^{14}\text{C}$  years BP (Iversen, 1954; Watts, 1980). While the magnitude, timing, and distribution of this event has had considerable attention in Europe itself since that time, it is only in the last two decades that the focus has expanded to include the extra-European presence of the signal. We define the Younger Dryas here as a chronozone (Mangerud *et al.*, 1974), because it is only through the  $^{14}\text{C}$  chronology that we may correlate vegetational and climatic events in various regions outside Europe. Unfortunately, a plateau may exist in the record at 10,000  $^{14}\text{C}$  years BP (Ammann and Lotter, 1989; Kromer and Becker, 1990), resulting in the same  $^{14}\text{C}$  ages over a stratigraphic interval due to variations in the proportion of  $^{14}\text{C}$  in the atmosphere. However, without tephra stratigraphy or other means of correlation we must rely on  $^{14}\text{C}$  chronology, and all chronology referenced in this paper is in radiocarbon years BP. The aims of the Global Younger Dryas? Working Group include the following:

- (1) To generate a critical assessment of the late-Pleistocene/Holocene transition pollen stratigraphy throughout various regions of the globe, excluding Europe, in order to determine whether or not a climatic reversal is present. This task is not an easy one for many reasons. Lack of close sampling intervals in many cores, lack of consistent regional pollen stratigraphies, and poor chronological control all contribute to the difficulty in addressing this question. Additionally, the palynologist faces the problem of translating a palynological change into vegetational change and then making a climatic interpretation. The participants of this working

group (Table 1) have made a large effort to include all existing  $^{14}\text{C}$  chronologies in regional reviews for the reader's benefit.

- (2) To produce new late-glacial pollen and glacial records of change which have good chronological control, preferably using accelerator mass spectrometry (AMS)  $^{14}\text{C}$  dating.
- (3) To provide an interdisciplinary focus from comparison of ice core stratigraphy, marine records, and glacial and pollen stratigraphy so as to improve our understanding of the mechanism and origin of this event.

TABLE 1. Participant Authors from IGCP 253 Working Group 7, Global Younger Dryas?

|   |
|---|
| R. Alley, Penn. State Univ., University Park, PA  |
| B. Ammann, Universitat Bern, Bern, Switzerland  |
| K.H. Anderson, Brown University, Providence, RI   |
| R.S. Anderson, Northern Arizona University, Flagstaff, Arizona                            |
| M. Arnold, Centre des Faibles Radioactivites, Gif-sur-Yvette, France                      |
| E. Bard, Université d'Aix-Marseille III, Marseille, France                                |
| P.B. Beaumont, McGregor Museum, Kimberley, South Africa                                   |
| H.J.B. Birks, University of Bergen, Bergen, Norway  |
| G. Chappellaz, Centre National de la Recherche Scientifique, Saint Martin d'Heres, France |
| R. Bonnefille, Laboratoire du Geologie du Quaternaire, Marseille, France                  |
| S. Bottema, Rijksuniversiteit, Groningen, The Netherlands                                 |
| G. Buchet, Laboratoire du Geologie du Quaternaire, Marseille, France                      |
| C. Clapperton, University of Aberdeen, Aberdeen, Scotland                                 |
| L. Cwynar, University of New Brunswick, Fredericton, New Brunswick, Canada                |
| O. Davis, University of Arizona, Tucson, Arizona  |
| P.T. Davis, Bentley College, Waltham, MA  |
| A. Del Genio, NASA/Goddard Institute for Space Studies, New York, NY                      |
| R. Drescher-Schneider, Universitat Bern, Bern, Switzerland                                |
| T. Goslar, Institute of Physics, Gliwice, Poland  |
| B. Hansen, University of Minnesota, Minneapolis, MN                                       |
| C.J. Heusser, Clinton Woods, Tuxedo, New York   |
| H. Hooghiemstra, University of Amsterdam, Amsterdam, The Netherlands                      |
| M. Icole, Lab du Geologie du Quaternaire, Marseille, France                               |
| S. Juggins, University College, London  |
| L. Keigwin, Woods Hole Oceanographic Institute, Woods Hole, MA                            |
| T. Kuc, Institute of Physics and Nuclear Techniques, Krakow, Poland                       |
| P. Kuhry, University of Alberta, Edmonton, Alberta, Canada                                |
| R. Lafonte, Lab de Geologie du Quaternaire, Marseille, France                             |
| G. Lang, Universitat Bern, Bern, Switzerland  |
| A. Levesque, University of New Brunswick, Fredericton, New Brunswick, Canada              |

TABLE 1. (Continued)

---

|  |
|--|
| B. Leyden, University of S. Florida, Tampa, FL   |
| A. Lotter, Universität Bern, Bern, Switzerland   |
| B. Maloney, The Queens University of Belfast, Belfast, Northern Ireland                                |
| D. Mann, University of Alaska, Fairbanks, Alaska   |
| V. Markgraf, University of Colorado, Boulder, Colorado   |
| R. Mathewes, Simon Fraser University, Burnaby, British Columbia, Canada                                |
| F. Mayle, University of New Brunswick, Fredericton, New Brunswick, Canada                              |
| M. McGlone, Landcare Research N. Zealand Ltd, Lincoln, New Zealand                                     |
| G. Osborne, University of Calgary, Calgary, Canada   |
| M. Pazdur, Institute of Physics, Gliwice, Poland   |
| D. Peteet, NASA/Goddard Institute for Space Studies, New York, NY                                      |
| M. Ralska-Jasiewiczowa, Polish Academy of Sciences, Krakow, Poland                                     |
| M. Reasoner, University of Alberta, Edmonton, Canada   |
| G. Rioulet, Lab de Geologie du Quaternaire, Marseille, France  |
| D. Rodbell, Byrd Polar Research Center, Columbus, OH   |
| M. Rossignol-Strick, Université P & M Curie, Paris, France   |
| K. Rozanski, International Atomic Energy Agency Vienna, Austria  |
| L. Scott, The University of the Orange Free State, Republic of South Africa                            |
| G. Seltzer, Byrd Polar Research Center, Columbus, OH   |
| D. Shafer, Northern Arizona University, Flagstaff, AZ  |
| L.C.K. Shane, University of Minnesota, Minneapolis, MN   |
| L. Sirkin, Adelphi University, Garden City, New York   |
| M. Steenkamp, University of Orange Free State, Bloemfontein, Republic of South Africa                  |
| K. Szeroczynska, Institute of Geological Sciences, Warszawa, Poland                                    |
| T. van der Hammen, University of Amsterdam, Amsterdam, The Netherlands                                 |
| T.R. Van Devender, Arizona-Sonora Desert Museum, Tucson, Arizona                                       |
| B. Van Geel, University of Amsterdam, Amsterdam, The Netherlands                                       |
| A. Walanus, Institute of Physics, Gliwice, Poland  |
| B. Wicik, Institute of Geography, Warszawa, Poland   |
| K. Wieckowski, Institute of Geography and Spatial Organization, Warszawa, Poland                       |
| Zielinski, G. Institute for Study of Earth, Oceans, and Space, University of New Hampshire, Durham, NH |

---

### Summary of Earlier Work

The late-glacial paleoecological history of the globe comprises a vast literature, which would be impossible to summarize in this review. Early reviews of Younger Dryas European pollen evidence (Watts, 1980) emphasized the dramatic changes in vegetation that characterized both coastal and inland areas. In areas outside Europe, few researchers correlated late-glacial fluctuations with the Allerød/Younger Dryas oscillation, but some exceptions exist, and are being investigated now with renewed interest. It was the discovery of a clear Younger Dryas (YD) signal in the Greenland ice cores in the last decade (Dansgaard *et al.*, 1989; Alley *et al.*, 1993; Taylor *et al.*, 1993) that sparked additional interest in the late-glacial climate. These high-resolution, multi-proxy records generated enhanced impetus to understand the cause of the dramatic shift in climate that characterized Greenland and Europe roughly 12,800–11,600 calendar years ago (ice-core chronology). The record of a YD methane signal in Antarctic ice cores (Chappellaz *et al.*, 1990) showed

the extent of the geographic record of the changes which were characteristic of this climatic interval.

In contrast to the European evidence for the YD, most North American data prior to the late 1980s was interpreted as indicative of uniform warming after ice retreat (Watts, 1983, Gaudreau and Webb, 1985). An exception to this pattern was the pollen and glacial evidence noted in the Great Lakes region of North America, which was related to the YD (Saarnisto, 1974; Bjorck, 1984; Shane, 1987) and the lithological and pollen data from sites in Atlantic maritime Canada (Mott *et al.*, 1986).

Taking a more global view, Broecker *et al.* (1985) and Rind *et al.* (1986) summarized the palynological, marine, and glacial evidence that supported a possible global YD expression, and provided a hypothesis and a model concerning its origin related to changes in ocean circulation. They noted that researchers in northeastern maritime Canada, Pacific northwestern North America, northern South America, southern South America, Russia, China, and Africa all had suggested that the late-glacial oscillation they recorded might be correlative with the YD (Rind *et al.*, 1986).

The importance of understanding the Younger Dryas event in its geographic extent, magnitude, and marine-ice-terrestrial connection has led to the formation of this working group (Table 2). Publications of this working group to date are listed in Table 3, and the global summary of the YD includes this research. There is, of course, considerable overlap with all other IGCP-253 working groups. For example, IGCP-253 Working Group No. 4, North Atlantic Seaboard Programme, has produced a volume of papers (Lowe, 1994) that addresses the late-glacial history of many regions surrounding the North Atlantic. In contrast to earlier views suggesting that the Younger Dryas was strictly a European event (i.e. Mercer, 1969), this set of papers includes summary data from Iceland, Nova Scotia, New Brunswick, Newfoundland, Quebec, Baffin Island, northern Labrador, and southern New England. Most of these regions now report a YD oscillation, except Quebec. Additions to the European YD are new summaries from the eastern Mediterranean region, suggesting that the oscillation characterized some sites there (Rossignol-Strick, *in press*; Bottema, *in press*).

Prior to the last decade, reliance on conventional dating greatly limited the strength of YD correlation. The recent technological advance of improved chronological resolution and precision with the AMS  $^{14}\text{C}$ -dating has resulted in a much stronger understanding of the timing of late-glacial events, and improved correlation with the European YD sites. The attempt to take an interdisciplinary view has also resulted in a much better appreciation of the complexity of the global climate system. Although many problems in correcting  $^{14}\text{C}$  ages to calendar years remain (Bard *et al.*, 1993), it is clear that the potential to link the response of the terrestrial system to ocean and atmospheric changes has improved during the last decade.

TABLE 2. Global Younger Dryas? Working Group Activities: Meetings

- **September, 1992**  
Global Younger Dryas? Symposium, International Palynological Meeting, Aix-en-Provence, France (organized by D. Peteet, N. Khotinsky, R. Mathewes). 150 attendees, with 11 papers presented. Abstracts published in IPC Program and Abstracts, 1992.
- **December 1992**  
Fall American Geophysical Union Mtg, San Francisco, California, special Symposium entitled High Latitude Paleoclimate Change (D. Peteet and V. Markgraf, presiding). 100 attendees, with 14 paper presentations representing seven countries. The focus of the meeting was the comparison of late-glacial ice core, marine, and terrestrial high latitude pollen and glacial records. The YD was targeted as a major item of interest for each of the various indices in widespread regions, ranging from Greenland, high latitude North Atlantic and North Pacific regions to South America and Antarctica. Abstracts of papers are published in EOS.
- **April 1993**  
PAGES (Past Global Climate Changes) and Lamont Climate Center sponsored a small Global Younger Dryas? Workshop at LDGO. 20 attendees, with seven paper presentations representing three countries. A very small, interdisciplinary group of researchers working on ice cores, glacial records, marine cores, pollen records, and atmospheric processes. Aim of the meeting was to produce a 10-pg document stating the state-of-art knowledge concerning the subject, gaps in our understanding, and recommendations for future research. Published in EOS 74: 587-589.
- **June 1993**  
Global Younger Dryas? Symposium, Winnipeg, Manitoba 50 + attendees, with seven paper presentations representing five countries. Focus of the meeting was to present papers concerning the presence or absence of an event correlative with the European Younger Dryas 11–10,000 <sup>14</sup>C years BP, within the context of the late-glacial interval. Abstracts published by J. Teller, Winnipeg, 1993.

TABLE 3. Publications: IGCP Global Younger Dryas?

- 8th Int. Palynological Congress Programs and Abstracts, 6–12 Sept. 1992
- Peteet, D. (ed.) (1993). Alley, R., Bond, G., Chappellaz, J., Clapperton, C., Del Genio, A., Keigwin, L. and Peteet, D. (1993). Global Younger Dryas?, *EOS*, 74 (50), 587–589.
- Peteet, D. (ed.) (1993). Global Younger Dryas?, *Quaternary Sci. Rev.*, 12, p.v. and list of papers
- Ammann, B., Birks, H.J.B., Drescher-Schneider, R., Juggins, S., Lang, G. and Lotter, A.F. (1993). Patterns of variation in late-glacial pollen stratigraphy along a Northwest–Southeast Transect Through Switzerland — a Numerical Analysis. *Quaternary Sci. Rev.*, 12, 277–286.
- Goslar, T., Kuc, T., Ralska-Jasiewiczowa, Rozanski, K., Arnold, M., Bard, E., van Geel, B., Pazdur, M.F., Szeroczynska, K., Wicik, B., Wieckowski, K. and Walanus, A. (1993). High-resolution lacustrine record of the late-glacial/Holocene transition in Central Europe. *Quaternary Sci. Rev.*, 12, 287–294.
- Mayle, F.E., Levesque, A.J. and Cwynar, L.C. (1993). *Alnus* as an indicator taxon of the Younger Dryas cooling in eastern North America. *Quaternary Sci. Rev.*, 12, 295–306.
- Shane, L.C.K. and Anderson, K.H. (1993). Intensity, gradients, and reversals in late glacial environmental change in east-central North America. *Quaternary Sci. Rev.*, 12, 307–320.
- Mathewes, R.W. (1993). Evidence for Younger Dryas-Age Cooling on the North Pacific coast of America. *Quaternary Sci. Rev.*, 12, 321–332.
- Kuhry, P., Hooghiemstra, H., Van Geel, B. and Van der Hammen, T. (1993). The El Abra stadial in the eastern Cordillera of Colombia (South America). *Quaternary Sci. Rev.*, 12, 333–344.
- Heusser, C.J. (1993). Late-glacial of Southern South America. *Quaternary Sci. Rev.*, 12, 345–350.
- Markgraf, V. (1993). Younger Dryas in southernmost South America. *Quaternary Sci. Rev.*, 12, 346–351.
- Peteet, D.M., Vogel, J.S., Nelson, D.E., Southon, J.R., Nickmann, R.J., and Heusser, L.E. (1990). Younger Dryas climatic reversal in northeastern USA? AMS ages for an old problem. *Quaternary Res.*, 33, 219–230.
- Peteet, D.M., Daniels, R., Heusser, L.E., Vogel, J.S., Southon, J.R. and Nelson, D.E. (1994). Wisconsinan late-glacial environmental change in southern New England: a regional synthesis. *J. Quaternary Sci.*, 9 (2), 151–154.

## EVIDENCE FOR EXTRA-EUROPEAN YOUNGER DRYAS

### North America

Recent attention to the late-glacial has resulted in the emerging re-examination of whether or not the YD is evident in various regions of North America. This summary of regional evidence focuses primarily on regions adjacent to the existing ice sheets, spanning the continent. While we focus on areas where this oscillation has been identified, other marginal ice sheet areas exist where the oscillation has not been recognized, i.e. Maine and Quebec.

### Eastern Canada

At the eastern margin of the Laurentide ice sheet, numerous late-glacial buried sequences, found more than half a century ago, indicate a sudden oscillation in climate (Mott, 1985; Mott *et al.*, 1986). For many years problems with conventional <sup>14</sup>C-dating precluded their precise correlation with the YD event, but the advent of AMS dating combined with very detailed macrofossil analysis has resulted in the strongest establishment of this correlation in North America

(Fig. 1). Colder climatic conditions resulted in the replacement of shrub-tundra (*Betula glandulosa*) with herb tundra (Cyperaceae, *Artemisia*, *Oxyria*) in central New Brunswick and northern Nova Scotia during the YD (Levesque *et al.*, 1993; Mayle *et al.*, 1993a,b; Mott, 1994).

In southern New Brunswick and central mainland Nova Scotia, closed forest (*Picea*) was replaced by shrub tundra (*Alnus crispa*, Cyperaceae, Poaceae) (Mayle *et al.*, 1993a,b; Cwynar *et al.*, 1994; Levesque *et al.*, 1994; Mott, 1994; Mayle and Cwynar, *in press*). The lake sediments are characterized by increased inorganic material from landscape erosion. In New Brunswick, diatom production was greatly diminished (Rawlence, 1988), and surface water temperature reconstructions derived from modern chironomid/temperature relationships indicate temperature declines of 6–7°C (Walker *et al.*, 1991; Wilson *et al.*, 1993) and at Pine Ridge Pond, New Brunswick, a decline of 12°C (Levesque *et al.*, 1994). In Newfoundland, accelerated erosion of surface soils is indicated by an increase in alkali in the sediment, and palynological data from eight sites show YD increases in various herb tundra species, e.g.




FIG. 1. Changes in percentage, concentration, and influx values for *Alnus crispa* pollen during the late-glacial and early Holocene in maritime Canada. The continuous horizontal lines mark the boundaries of the Younger Dryas as determined from changes in percent LOI (From Mayle *et al.*, 1993).

Ericales and Cyperaceae (Anderson and MacPherson, 1994).

In contrast to this pattern of coastal Canadian response to the Younger Dryas cooling, palynological records from Quebec do not appear to record a significant signal (Richard, 1994) during the YD, but a cold episode in the Holocene is noted at 9.5  $^{14}\text{C}$  years BP.

#### Southern New England

Southern New England shows a strong regional pattern of late-glacial palynological change (Peteet, 1987), that is apparent in over twenty lake records and has been recognized since the 1950s (Leopold, 1956). However, since that time the changes have been interpreted as indicating unidirectional warming (e.g. Davis, 1967; Watts, 1983; Gaudreau and Webb, 1985). Whether or not this palynological change represented a regional vegetational change was debated for many years, which precluded its definition as a YD correlative. Macrofossil analysis and pollen influx from three of these lakes have resulted in the documentation

of the presence of temperate species in southern New England prior to the YD cooling, while  $^{14}\text{C}$  AMS dates provide a chronology for the timing of the YD response (Peteet *et al.*, 1990, 1993, 1994). Regionally, organic sedimentation began abruptly approximately 12,400 years BP, when the ice had retreated enough for the lake basins to form. Fluctuations in a mixed coniferous-deciduous forest (*Picea*, *Abies*, *Larix*, *Pinus strobus*, *Quercus*, *Fraxinus*) culminate about 11,000 years BP with a rapid disappearance of the thermophilous trees and a marked increase in the boreal indicators (*Picea*, *Abies*, *Larix*, *Betula papyrifera*, *Alnus*) as climate cooled (Fig. 2). This palynological signal is supported by dominance of the boreal macrofossils and the disappearance of *Pinus strobus* needles. We infer this vegetational change to be indicative of an approximate July cooling of 3–4°C (Peteet *et al.*, 1990, 1993). The sharp return to warmer conditions at 10,000  $^{14}\text{C}$  BP was marked by the dominance of *Pinus strobus* and loss of the boreal component.

Close-interval analyses of a pollen and macrofossil late-glacial record from the Hudson Highlands of




FIG. 2. Pollen percentage diagram of selected types, 8–12 m depth, Linsley Pond, Conn. (top). Macrofossil diagram from 10 to 12 m depth, Linsley Pond, Conn. Samples represent material from 50 cms of sediment at 5 cm intervals. *Pinus strobus*, *Abies*, and *Larix* macrofossils are needles, *Betula papyrifera* remains include cone scales and seeds, and the remainder denote seeds or fruits (bottom) (from Peteet *et al.*, 1993).




FIG. 3. Map showing intensity gradient in the *Picea* recurrence. Only sites with interpretable records are included. Numbers correspond to Table 1 in Shane *et al.* (1993). Open circles are sites with no *Picea* fluctuation in the pollen record; closed triangles are sites with a strong *Picea* pollen fluctuation (increase from a low point of over ca. 7%); open triangles are sites with a minor *Picea* pollen fluctuation (from Shane and Anderson, 1993).

southeastern New York provide an additional record of YD cooling with a dominance of *Abies* and *Alnus* 11–10,000 years BP (Maenza-Gmelch, 1994). These results contrast with the apparent lack of a YD signal in northern New England, where no climatic oscillation has been identified (Jacobson *et al.*, 1987).

#### Midwestern U.S.A.

From scrutiny of more than 25 conventional radiocarbon-dated pollen sites in the Allegheny Plateau (east) and Till Plains (west) of the Great Lakes region, Shane (1987) and Shane and Anderson (1993) have noted a cold oscillation equivalent to the YD in the western region. This cooling is defined by the rapid recurrence of *Picea* at the close of the late-glacial in the Till Plains sites (Fig. 3). Transfer function estimates show a January temperature drop of  $-5$  to  $-2^{\circ}\text{C}$  and a July temperature drop of  $1$ – $2^{\circ}\text{C}$ , with a precipitation increase from the Allerød. Whether or not the climate reversal represents a direct result of glacial and meltwater changes in the Great Lakes region or is due to extra-regional forcing is not yet resolved (Shane and Anderson, 1993). Future attention to AMS dating is needed to refine the chronology.

#### Pacific Northwest Coast

The Pacific Northwest record of palynological change from the Olympic Peninsula northwestward to coastal Alaska exhibits climatic oscillations during the late-glacial (Heusser, 1960). Increases in *Tsuga mertensiana* are the most notable link in more than ten sites to a Younger Dryas cooling along the British Columbia coastline (Mathewes, 1973; Hebda, 1983; Mathewes, 1993). The pollen percentage increase is small (3–5%), but mountain hemlock is notable for its very low pollen production, and a percentage increase is interpreted as a definite forest change (Fig. 4). This late-glacial increase from many sites suggests a decline in temperature and increased moisture in comparison with preceding centuries. AMS studies in progress will further constrain the timing of this oscillation.

To the north, one site in Glacier Bay, southeastern Alaska, shows an expansion of tundra elements concurrent with increased inorganic sediments between 10,600 and 9990 years BP and has thus been interpreted as a possible YD oscillation (Engstrom *et al.*, 1990).

Northwestward, a suite of coastal sections from Kodiak Island exhibits an abrupt lithological oscillation from 10,800 to 10,000 AMS  $^{14}\text{C}$  years BP (Peteet




FIG. 4. Generalized pollen diagram for Marion Lake, British Columbia, with climate interpretation and geologic-climate units of southwestern BC that are associated with cooling. YD = Younger Dryas chronozone (11–10 ka BP) with high mountain hemlock (*Tsuga mertensiana*) pollen; Sumas = climax of Sumas Stage ice readvance into Fraser Lowland around 11.3 ka BP (from Mathewes, 1993).

and Mann, 1994). Palynological and macrofossil data completed from one site support the change as a rapid and dramatic cooling, with loss of Polypodiaceae and dominance of *Empetrum* and *Artemisia*. The return to warm conditions at 10,000  $^{14}\text{C}$  years BP is very abrupt.

The pattern, magnitude, and precise timing of these vegetational and climatic changes throughout North America deserves additional attention. Problems with radiocarbon plateaus at 12,700 and 10,000  $^{14}\text{C}$  years BP during the late-glacial to postglacial transition (Ammann and Lotter, 1989; Kromer and Becker, 1990) complicate the estimation of rates of change. Many questions remain as to the geographic extent of the cooling and why significant gaps appear to exist (i.e. Quebec, northern New England). The extent of the oscillation as one moves southward is also an important question, as references have been made to its possible occurrence in Kentucky (Wilkins *et al.*, 1991). Comparison of the terrestrial data with ice core and marine records indicates the importance of consideration of migration lags, thresholds, and separation of temperature and moisture signals. Evidence of a possible North Pacific YD equivalent suggests that the origin of the terrestrial cooling may not have been restricted to the North Atlantic, or that the atmospheric response was very rapid and widespread (Peteet, 1993a,b).

#### Central America

Leyden (*in press*) has summarized late-glacial evidence from several wetlands in Central America.

Two of these records, La Chonta bog area, Costa Rica (Hooghiemstra *et al.*, 1992) and Lake Quexil, Guatemala (Deevey *et al.*, 1983; Leyden, 1984; Leyden *et al.*, 1993) exhibit late-glacial reversals, estimated to have begun 12–11,000 years BP and ending before 10,300 years BP, but dating is not precise enough to establish a YD correlative event. Lake La Yeguada (Piperno *et al.*, 1990; Bush *et al.*, 1992) from lowland Panama, in contrast, does not show a clear late-glacial reversal.

#### South America

##### Colombia

In 1959, Van der Hammen and Gonzalez published a record of late-glacial climatic reversal in Colombian montane cores of the Eastern Cordillera. They named the warming the Guantiva interstadial, which was closely followed by the El Abra stadial (colder), and then the Holocene. Continued investigation of many sites (e.g. Fig. 5) by a variety of authors has shown this to be a regional palynological reversal (see Kuhry *et al.*, 1993), occurring approximately 11–10,000  $^{14}\text{C}$  years BP, correlative with the Allerod-Younger Dryas oscillation (Kuhry *et al.*, 1993; Van der Hammen and Hooghiemstra, *in press*). Colder climatic conditions during the El Abra stadial caused a 600 m lowering of the upper Andean forest limit and a cooling of 4°C compared to the present. Available conventional dating shows the transition from the Guantiva interstadial to the El Abra stadial to be  $11,210 \pm 90$  and  $10,820 \pm 60$ , and warming may have started shortly after  $10,380 \pm 90$  (Kuhry *et al.*, 1993). AMS dating is presently in progress to establish more closely the timing of the event (Van der Hammen and Hooghiemstra, *in press*).

In the Sierra Nevada del Cocuy, a glacial advance correlative with the El Abra stadial was named the Bocatoma stade (Gonzalez *et al.*, 1965). The ELA lowering was about 700 m.

In contrast to the correlation of the Colombian pollen and glacial stratigraphy with the YD event, Heine (1993) suggests that the evidence for a YD oscillation is as yet inconclusive, and that more dating is needed to establish this event in the tropical Andes.

##### Ecuador and Peru

According to a review by Hansen (*in press*) of the published palynological records that exist from Ecuador, none exhibit a late-glacial oscillation and all lack radiocarbon dating delineating the YD. In Peru, pollen data from four sites suggest one and possibly two late-glacial climatic oscillations but only one site, Laguna Junin, has dates which bracket the short-term climate change, and it appears to be between 12 and 11,000 years BP. No AMS dates are as yet available.

##### Chile and Argentina

Whether or not a late-glacial climatic reversal characterized southernmost South America has been




FIG. 5. Simplified pollen diagram from the upper part of the Laguna de Fuquene II core (Van Geel and Van der Hammen, 1973). Additional radiocarbon date from the Laguna de Fuquene III core (From Kuhry *et al.*, 1993).

controversial for almost a decade. Two recent reviews present both sides of the argument (Heusser, 1993; Markgraf 1993). While some records clearly do show an oscillation between about 11 and 10,000 years ago, such as Alerce and Rucanancu (Heusser, 1989, 1993), others such as Rio Caunahue (Heusser, 1981; Markgraf, 1991) do not. Markgraf (1993) ascribes some of the late-glacial variability in the records to the effects of fire. No AMS dates are available as yet to refine the chronologies.

#### Potential Glacial YD Evidence — the Americas

Osborne *et al.* (*in press*) present the late-glacial alpine evidence for a possible Younger Dryas equivalent. The best examples are in the Canadian (Crowfoot moraine; Osborne *et al.*, *in press*) and American Rockies (Triple Lakes, Colorado; Davis and Osborne, 1987), the Ecuadorian (Reschreiter moraine; Clapperton, 1990), Peruvian (Cordillera Oriental and Cordillera Blanca; Rodbell, 1991, 1993), and Bolivian Andes, and the Patagonian ice cap. Most are not dated well enough to correlate with the YD, but two readvances fall within the YD interval: an advance of the Crowfoot Glacier in Canada, and the Reschreiter Glacier in Ecuador.

Heine (1993) suggests alternative explanations for the glacial chronologies from Ecuador.

#### Equatorial and East Africa

Coetzee (1967) suggested that a late-glacial palynological oscillation from Mt Kenya was related to moisture. New data from Burundi, equatorial Arica (Bonnefille *et al.*, *in press*) with AMS dates suggest that a vegetational oscillation did characterize the late-glacial. However, additional AMS-dated sites are needed to establish a regional climatic oscillation and to correlate with a YD signal.

#### South Africa

Scott *et al.* (*in press*) review the late-glacial evidence for a climatic oscillation in South Africa. From a total of 15 sites, they find that less than half have adequate resolution or dating to determine whether or not an oscillation exists. None of the palynological sequences in the interior so far suggest a Younger Dryas equivalent.



### South East Asia

Maloney (*in press*) reviews the available evidence from Southeast Asia concerning the late-glacial. He notes that the clearest evidence for a possible climatic oscillation is found in west Java and north Sumatra, but both sites need better dating.

### New Zealand

The available late-glacial pollen evidence for New Zealand, summarized by McGlone (*in press*), suggests that no significant reversal of the trend towards afforestation occurred after glaciation. However, many of the sites have poor sampling and chronological resolution. In contrast to the lack of palynological evidence for a YD in New Zealand, Denton and Hendy (1994) recently demonstrated a glacial advance of the Franz Josef Glacier to be a YD correlative.

## SUMMARY AND FUTURE RESEARCH

A summary map of the global palynological evidence for the YD is presented in Fig. 6. One of the major contributions of this working group has been to establish the clear evidence for the YD outside Europe,

in eastern North America. It is quite evident from the macrofossil evidence and the AMS  $^{14}\text{C}$ -dating that vegetation responds very quickly to climatic change, both at the onset and the close of the YD. The probable extent of the YD in the midwestern U.S.A. and along the North Pacific coast is indicated by a palynological oscillation at many sites, but they are not characterized by firm chronology. The possible occurrences in Glacier Bay, Alaska, and Kodiak, Alaska await further palynological investigations.

The most convincing evidence outside North America is from Colombia, South America, where many sites show an oscillation but await clear AMS  $^{14}\text{C}$  dating. Controversial regions where some sites seem to have an oscillation and others do not are Central America, southern South America, and the eastern Mediterranean region. A clear absence of a YD signal is not yet confirmed in any area, because high-resolution AMS-dated sites are rare, and none have been cited to prove that an oscillation is lacking during this interval. Regions where existing data suggest that no palynological oscillation took place include Ecuador and Peru, South Africa, and New Zealand, but in some of these same areas (New Zealand, possibly Peru) glacial advances do indicate a YD correlative.




FIG. 6. Global map of distribution of palynological evidence for the Younger Dryas cooling 11–10,000 yr BP. Where clear palynological evidence exists for a climatic oscillation in three or more sites, with AMS  $^{14}\text{C}$  dating, Y denotes a Yes YD (e.g. maritime eastern Canada). For regions where three or more sites show a palynological oscillation, but it is not yet AMS-dated, a PR is designated for Probable YD (i.e. Colombia). For sites where one or two oscillations exist, a PO denotes Possible YD (e.g. southern Alaska). For sites where some sites show an oscillation and others do not, a C denotes Controversial. Finally, where evidence for a YD is lacking, a N? is given for No? YD, unless three or more AMS sites have been investigated, in which case the designation would be N for No YD.

Examination of Fig. 6 reveals large areas of the continents where no designation for a YD can be made. Some of these continental areas were ice-covered during the YD (e.g. large portions of Canada) but in other regions lack of investigation or translation of literature into English (the former USSR) makes the areas difficult to assess. In particular, little appears to be known of India, Africa, large areas of S. America, and China. It is our hope that renewed interest in the YD will stimulate further pollen stratigraphic investigations in these regions in the future.

Through high-resolution sampling and dating in many regions of the globe, we eventually hope to be in a position to state with confidence whether or not the YD was a global event, and whether or not there were leads and lags in the expression of both vegetational and climatic change. By focused attention to interpretation of vegetational records, we hope to refine our estimates of temperature or moisture change. Careful analysis of the timing of the YD event in ice core, marine, and terrestrial records will help to expand the pattern of its geographic expression, and ultimately, may help us understand the mechanism of this dramatic climatic change.

## ACKNOWLEDGEMENTS

Many contributors to the *Global Younger Dryas?* have made the subproject stimulating to all participants. In particular, Jan Lundquist (Project Leader of IGCP-253) and Matti Saarnisto (Secretary of IGCP-253) conceived the project and supported its international scope. Discussions with other subproject and regional leaders broadened the perspective. Jose Mendoza and Lily Del Valle assisted in drafting figures. Thanks are due to M. Walker and H.H. Birks for thoughtful reviews of the manuscript. The author is indebted to PAGES for sponsoring the Workshop at Lamont-Doherty Earth Observatory. This research was also supported by NASA and NIGEC Northeast Regional Center at Harvard. This is Lamont Doherty Earth Observatory contribution, no. 5370.

## REFERENCES

- Alley, R.B., Meese, D.A., Shuman, C.A. *et al.* (1993). Abrupt increase in Greenland snow accumulation at the end of the Younger Dryas event. *Nature*, **362**, 527–529.
- Ammann, B. and Lotter, A.F. (1989). Late-glacial radiocarbon and palynostratigraphy on the Swiss Plateau. *Boreas*, **18**, 109–126.
- Anderson, T.W. and MacPherson, J.B. (1994). Wisconsinan late-glacial environmental change in Newfoundland: a regional review. *J. Quaternary Sci.*, **9**, 171–178.
- Bard, E., Arnold, M., Fairbanks, R. and Hamelin, B. (1993).  $^{230}\text{Th}$ - $^{234}\text{U}$  and  $^{14}\text{C}$  ages obtained by mass spectrometry on corals. *Radiocarbon*, **35**, 191–199.
- Björck, B. (1984). Deglaciation chronology and revegetation in northwestern Ontario. *Can. J. E. Sci.*, **22**, 850–871.
- Bonnefille, R., Riollet, G., Buchet, G., Arnold, M., Icole, M. and Lafont, R. (*in press*) Rusaka, a high resolution pollen record for the glacial-interglacial transition in equatorial Africa. *Quaternary Science Reviews*.
- Bottema, S. (*in press*). The Younger Dryas in the eastern Mediterranean. *Quaternary Sci. Rev.*.
- Broecker, W., Peteet, D. and Rind, B. (1985). Does the ocean-atmosphere system have more than one stable mode of operation? *Nature*, **315**, 21–26.
- Bush, M.B., Piperno, D.R., Colinvaux, P.A., De Oliveira, P.E., Kriesek, L.A., Miller, M.C. and Rowe, W.E. (1992). A 14,300-yr paleoecological profile of a lowland tropical lake in Panama. *Ecol. Monographs*, **62**, 251–275.
- Chappellaz, J., Barnola, J.M., Raynaud, D., Korotkevitch, Y.S. and Lorius, C. (1990). Atmospheric  $\text{CH}_4$  record over the last climatic cycle revealed by the Vostok ice core. *Nature*, **345**, 127–131.
- Clapperton, C.M. (1990). Glacial and volcanic geomorphology of the Chimborazo-Carihuairazo massif, Ecuadorian Andes. *Trans. Royal Soc. Edinburgh*, **81**, 91–116.
- Coetzee, J.A. (1967). Pollen analytical studies in East and Southern Africa. *Palaeoecol. of Africa*, **3**, 1–146.
- Cwynar, L.C., Levesque, A.J., Mayle, F.E. and Walker, I.R. (1994). Wisconsinan late-glacial environmental change in New Brunswick: a regional synthesis. *J. Quaternary Sci.*, **9**, 161–164.
- Dansgaard, W., White, J.W. and Johnsen, S.J. (1989). The abrupt termination of the Younger Dryas climatic event. *Nature*, **339**, 532–534.
- Davis, M.B. (1967). Late-glacial climate in northern United States: a comparison of New England and Great Lakes region. In *Quaternary Paleoeecology*, Cushing, E.J. and Wright, H.E., eds, pp. 11–44.
- Davis, P.T. and Osborne, G. (1987). Age of pre-Neoglacial cirque moraines in the central North American Cordillera. *Geographie Physique et Quaternaire*, **41**, 365–375.
- Deevey, E.S., Brenner, M. and Binford, M.W. (1983). Paleolimnology of the Peten lake district, Guatemala, III. Late Pleistocene and Gamblian environments of the Maya area. *Hydrobiologia*, **103**, 211–216.
- Denton, G.H. and Hendy, C.H. (1994). Younger Dryas age advance of Franz Josef Glacier in the Southern Alps of New Zealand. *Science*, **264**, 1434–1437.
- Engstrom, D.R., Hansen, B.C.S. and Wright, H.E. Jr (1990). A possible Younger Dryas record in southeastern Alaska. *Science*, **250**, 1383–1385.
- Gaudreau, D.C. and Webb, T. III (1985). Late-Quaternary pollen stratigraphy and isochron maps for the northeastern United States. In: Bryant, V. and Holloway, R.G. (eds). *Pollen Records of Late-Quaternary North American Sediments*, pp. 247–280. Amer. Assoc. of Stratigraphic Palynologists Foundation, Dallas, Texas.
- Gonzalez, E., Van der Hammen, T. and Flint, R.F. (1965). Late Quaternary glacial and vegetational sequence in Valle de Lagunillas, Sierra Nevada del Cocuy, Colombia. *Leidse Geologische Mededelingen*, **32**, 157–182.
- Hansen, B.C.S. (*in press*). A review of late-glacial pollen records from Ecuador and Peru with reference to the Younger Dryas event. *Quaternary Sci. Rev.*.
- Hebda, R.J. (1983). Late-glacial and postglacial vegetation history at Bear Cove Bog, northeast Vancouver Island, BC. *Can. J. Bot.*, **61**, 3172–3192.
- Heine, J.T. (1993). A reevaluation of the evidence for a Younger Dryas climatic reversal in the tropical Andes. *Quaternary Sci. Rev.*, **12**, 769–779.
- Heusser, C.J. (1960). *Late Pleistocene Environments of North Pacific North America*. Amer. Geogr. Soc. Sp. Publ. 35. New York.
- Heusser, C.J. (1981). Palynology of the last interglacial-glacial cycle in midlatitudes of southern Chile. *Quaternary Res.*, **16**, 293–321.
- Heusser, C.J. (1989). Late Quaternary and climate of southern Tierra del Fuego. *Quaternary Res.*, **31**, 396–406.
- Heusser, C.J. (1993). Late-glacial of Southern South America. *Quaternary Sci. Rev.*, **12**, 345–350.
- Hooghiemstra, H., Cleef, A.M., Noldus, G.W., and Kappelle, M. (1992). Upper Quaternary vegetation dynamics and paleoclimatology of the La Chonta bog area (Cordillera de Talamanca, Costa Rica). *J. Quaternary Sci.*, **7**, 205–225.
- Iversen, J. (1954). The lateglacial flora of Denmark and its relation to climate and soil. *Danm. Geol. Unders. Ser. II.*, **80**, 87–119.
- Jacobson, G.L., Webb, T. III. and Grimm, E. (1987). Patterns and rates of vegetational change during the deglaciation of eastern North America. In: Ruddiman, W.F. and Wright, H.E. Jr (eds), *North America and Adjacent Oceans during the Last Deglaciation*, pp. 277–288, Geology of North America, Vol. K-3, Boulder, CO.
- Jensen, K. (1938). Some west Baltic pollen diagrams. *Quartar*, **1**, 124–139.
- Kromer, B. and Becker, B. (1990). Tree ring  $^{14}\text{C}$  calibration at

- 10,000 BP. In: Bard, E. and Broecker, W.S. (eds). *The Last Deglaciation: Absolute and Radiocarbon Chronologies*, NATO ASI Series, Vol. 12, Springer-Verlag, New York.
- Kuhry, P., Hooghiemstra, H., Van Geel, B. and Van der Hammen, T. (1993). The El Abra Stadial in the Eastern Cordillera of Colombia (South America). *Quaternary Sci. Rev.*, **12**, 333–344.
- Leopold, E.B. (1956). Two late-glacial deposits in southern Connecticut. *Proc. National Acad. Sci.*, **42**, 863–867.
- Levesque, A.J., Mayle, F.E., Walker, I.R. and Cwynar, L.C. (1993). A previously unrecognized late-glacial cold event in eastern North America. *Nature*, **361**, 623–626.
- Levesque, A.J., Cwynar, L.C. and Walker, I.R. (1994). A multiproxy investigation of late-glacial climate and vegetation change at Pine Ridge Pond, Southwest New Brunswick, Canada. *Quaternary Res.*, **42**, 316–327.
- Leyden, B.W. (1984). Guatemalan forest synthesis after Pleistocene aridity. *Proc. National Acad. Sci. U.S.A.*, **81**, 4856–4859.
- Leyden, B.W. (in press). Evidence of the Younger Dryas in Central America. *Quaternary Sci. Rev.*
- Leyden, B.W., Brenner, M., Hodell, D.A. and Curtis, J.H. (1993). Late Pleistocene climate in the Central American lowlands. In: Swart, P.K., Lohmann, K.C., McKenzie, J. and Savin, S. (eds.), *Climate Change in Continental Isotopic Records*, pp. 165–178. Geophysical Monograph 78, American Geophysical Union, Washington, DC.
- Lowe, J.J. (1994). (Editor). North Atlantic Seaboard Programme IGCP-253, Climate Changes in Areas Adjacent to the North Atlantic during the last Glacial-Interglacial Transition. *J. Quaternary Sci.*, **9** (2), 95–198.
- Maenza-Gmelch, T. (1994). Paleoecology of the Late-glacial/Holocene transition in the Hudson Highlands, NY, USA. AMQUA 1994 Abstract, p. 124.
- Maloney, B.K. (in press). Evidence for the Younger Dryas event in Southeast Asia. *Quaternary Sci. Rev.*
- Mangerud, J., Anderson, S.T., Berglund, B.E. and Donner, J.J. (1974). Quaternary stratigraphy of Norden, a proposal for terminology and classification. *Boreas*, **3**, 109–127.
- Markgraf, V. (1991). Younger Dryas in southern South America? *Boreas*, **20**, 63–69.
- Markgraf, V. (1993). Younger Dryas in southernmost South America — an update. *Quaternary Sci. Rev.*, **12** (5), 351–355.
- Mathewes, R.W. (1973). A palynological study of postglacial vegetation changes in the University Research Forest, southwestern British Columbia. *Can. J. Botany*, **51**, 2085–3103.
- Mathewes, R.W. (1993). Evidence for Younger Dryas age cooling on the North Pacific Coast of America. *Quaternary Sci. Rev.*, **12**, 321–332.
- Mayle, F.E. and Cwynar, L.C. (in press). A review of multi-proxy data for the Younger Dryas in maritime eastern Canada. *Quaternary Sci. Rev.*
- Mayle, F.E., Levesque, A.J. and Cwynar, L.C. (1993a). Accelerator mass spectrometry ages for the Younger Dryas event in Atlantic Canada. *Quaternary Res.*, **39**, 355–360.
- Mayle, F.E., Levesque, A.J. and Cwynar, L.C. (1993b). *Alnus* as an indicator taxon of the Younger Dryas cooling in eastern North America. *Quaternary Sci. Rev.*, **12**, 295–305.
- McGlone, M.S. (in press). Landscape and vegetation change during the Younger Dryas interval in New Zealand. *Quaternary Sci. Rev.*
- Mercer, J.H. (1969). The Allerød oscillation: a European climate anomaly? *Arc. Alp. Res.*, **1**, 227–234.
- Mott, R.G. (1985). Late-glacial change in the maritime provinces. In: Harington, C.R. (ed.) *Climatic Change in Canada* 5, pp. 281–300.
- Mott, R.G. (1994). Wisconsinan late-glacial environmental change in Nova Scotia: a regional synthesis. *J. Quaternary Sci.*, **9**, 155–160.
- Mott, R.G., Grant, D.R., Stea, R. and Ochietti, S. (1986). Late-glacial climatic oscillation in Atlantic Canada equivalent to the Allerød-Younger Dryas event. *Nature*, **323**, 247–250.
- Osborne, G., Clapperton, C., Davis, P.T., Reasoner, M., Rodbell, D.T., Seltzer, G.O. and Zielinski, G. (in press). Potential glacial evidence for the Younger Dryas event in the cordillera of North and South America. *Quaternary Sci. Rev.*
- Peteet, D.M. (1987). Younger Dryas in North America — modeling, data analysis, and re-evaluation. In: Berger, W.H. and Labeyrie, L.D. (eds.), *Abrupt Climatic Change*, pp. 185–193. Reidel, Dordrecht.
- Peteet, D.M. (Ed.). (1993a). Global Younger Dryas? *Quaternary Sci. Rev.*, **12**(5).
- Peteet, D.M. (Ed.). (1993b). Global Younger Dryas? *EOS*, **74**(50), 587–589.
- Peteet, D.M. and Mann, D.H. (1994). Late-glacial vegetational, tephra, and climatic history of southwestern Kodiak Island, Alaska. *Ecoscience*, **1**(3), 255–267.
- Peteet, D.M., Vogel, J.S., Nelson, D.E., Southon, J.R., Nickmann, R.J. and Heusser, L.E. (1990). Younger Dryas reversals in northeastern USA? AMS ages for an old problem. *Quaternary Res.*, **33**, 219–230.
- Peteet, D.M., Daniels, R.A., Heusser, L.E., Vogel, J.S., Southon, J.R. and Nelson, D.E. (1993). Late-glacial pollen, macrofossils, and fish remains in the northeastern USA — the Younger Dryas oscillation. *Quaternary Sci. Rev.*, **2**, 597–612.
- Peteet, D.M., Daniels, R., Heusser, L.E., Vogel, J.S., Southon, J.R. and Nelson, D.E. (1994). Wisconsinan late-glacial environmental change in southern New England: a regional synthesis. *Quaternary Sci.*, **9**, 151–154.
- Piperno, D.R., Bush, M.B. and Colinvaux, P.A. (1990). Paleo-environments and human settlements in late-glacial Panama. *Quaternary Res.*, **33**, 108–116.
- Rawlence, D.J. (1988). The post-glacial diatom history of Splan Lake, New Brunswick. *J. Paleolimn.*, **1**, 51–60.
- Richard, P.J.H. (1994). Wisconsinan late-glacial environmental change in Quebec: a regional synthesis. *J. Quaternary Sci.*, **9**, 165–170.
- Rind, D., Peteet, D., Broecker, W., McIntyre, A. and Ruddiman, W. (1986). The impact of cold North Atlantic sea surface temperatures on climate: implications for the Younger Dryas cooling (II-10k). *Climate Dynamics*, **1**, 3–33.
- Rodbell, D.T. (1991). Late Quaternary glaciation and climatic change in the northern Peruvian Andes. Ph.D. dissertation, University of Colorado.
- Rodbell, D.T. (1993). Subdivision of late Pleistocene moraines in the Cordillera Blanca, Peru based on rock weathering features, soils, and radiocarbon dates. *Quaternary Res.*, **39**, 133–143.
- Rosignol-Strick, M. (in press). Late Quaternary climate in the Eastern Mediterranean region. *Quaternary Sci. Rev.*
- Saarnisto, M. (1974). The deglaciation history of the Lake Superior region and its climatic implications. *Quaternary Res.*, **4**, 316–339.
- Scott, L., Steenkamp, M. and Beaumont, P.B. (in press). Paleoenvironmental conditions in South Africa at the Pleistocene-Holocene transition. *Quaternary Sci. Rev.*
- Shane, L.C.K. (1987). Late-glacial vegetational and climatic history of the Allegheny Plateau and the Till Plains of Ohio and Indiana, U.S.A. *Boreas*, **16**, 1–20.
- Shane, L.C.K. and Anderson, K.H. (1993). Intensity, gradients, and reversals in late-glacial environmental change in east-central North America. *Quaternary Sci. Rev.*, **12**, 307–320.
- Taylor, K.C., Lamorey, G.W. et al. (1993). The “flickering switch” of late Pleistocene climatic change. *Nature*, **361**, 432–436.
- Van der Hammen, T. and Gonzalez, E. (1959). Historia de clima y vegetación del Paramo de Palacio (Cordillera Oriental, Dept. Cundinamarca). Official report of the Servicio Geológico Nacional, Bogotá; 19 pg.
- Van der Hammen, T. and Hooghiemstra, H. (in press). The El Abra stadial, a Younger Dryas equivalent in Colombia. *Quaternary Sci. Rev.*
- Van Geel, B. and Van der Hammen, T. (1973). Upper Quaternary vegetational and climatic sequence of the Fuquene area (Eastern Cordillera, Colombia). *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, **14**, 9–92.
- Walker, I.R., Mott, R.J. and Smol, J.P. (1991). Allerød-Younger Dryas lake temperatures from midge fossils in Atlantic Canada. *Science*, **253**, 1010–1012.
- Watts, W.A. (1980). Regional variations in the response of vegetation of late-glacial climate events in Europe. In: Lowe, J.J., Gray, J.M., and Robinson, J.E. (eds.), *The Late-Glacial of Northwest Europe*, pp. 1–22. Pergamon, New York.
- Watts, W.A. (1983). Vegetational history of the eastern United States 25000 to 10000 years ago. In: Porter, S.C. (ed.), *Late Quaternary Environments of the U.S.*, Vol. 2, pp. 294–310. University of Minnesota.
- Wilkins, G.R., Delcourt, P.A., Delcourt, H.R., Harrison, F.W. and Turner, M.R. (1991). Paleocology of central Kentucky since the last glacial maximum. *Quaternary Res.*, **36**, 224–239.
- Wilson, S.E., Walker, I.R., Mott, R.J. and Smol, J.P. (1993). Climate and limnological changes associated with the Younger Dryas in Atlantic Canada. *Climate Dynamics*, **8**, 177–187.

**APPENDIX: PAPERS FOR NEXT GLOBAL  
YOUNGER DRYAS? EDITION OF *QUATERNARY  
SCIENCE REVIEWS***

- Bonnefille, R., Riollet, G., Buchet, G., Arnold, M., Icole, M. and Lafont, R. Rusaka, a High Resolution Pollen Record for the Glacial/Interglacial Transition in Equatorial Africa.
- Bottema, S. The Younger Dryas in the eastern Mediterranean.
- Leyden, B.W. Evidence of the Younger Dryas in Central America.
- Hansen, B.C.S. A review of late-glacial pollen records from Ecuador and Peru with reference to the Younger Dryas event.
- Maloney, B.K. Evidence for the Younger Dryas event in Southeast Asia.
- Mayle, F.E. and Cwynar, L.C. A review of multi-proxy data for the Younger Dryas in maritime eastern Canada.
- McGlone, M.S. Landscape and vegetation change during the Younger Dryas interval in New Zealand.
- Osborne, G., Clapperton, C., Davis, P.T., Reasoner, M., Rodbell, D.T., Seltzer, G.O., and Zielinski, G. Potential glacial evidence for the Younger Dryas event in the cordillera of North and South America.
- Rosignol-Strick, M. Late Quaternary climate in the Eastern Mediterranean region.
- Scott, L. Steenkamp, M., and Beaumont, P.B. (1993). Paleoenvironmental conditions in South Africa at the Pleistocene-Holocene transition.
- Van der Hammen, T., and Hooghiemstra, H. The El Abra stadial, a Younger Dryas equivalent in Colombia.